



TITLE:

# Sufficient Conditions for Univalent Functions

AUTHOR(S):

NUNOKAWA, Mamoru

---

CITATION:

NUNOKAWA, Mamoru. Sufficient Conditions for Univalent Functions. 数理解析研究所講究録 1993, 821: 1-3

ISSUE DATE:

1993-02

URL:

<http://hdl.handle.net/2433/83199>

RIGHT:

# Sufficient Conditions for Univalent Functions

By Mamoru NUNOKAWA

Department of Mathematics, University of Gunma

(布川 護, 群馬大学)

Let  $A(p)$  denote the class of functions of the form

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n \quad (p \in \mathbb{N} = \{1, 2, 3, \dots\})$$

which are analytic in the unit disk  $U = \{z : |z| < 1\}$ .

Ozaki, Ono and Umezawa [5] proved the following result.

Theorem A. If  $f(z) \in A(1)$  and  $|f''(z)| < 1$  in  $U$ , then  $f(z)$  is univalent in  $U$ .

Recently, Nunokawa, Kwon and Cho [2] have proved the following result.

Theorem B. Let  $p \geq 2$ . If  $f(z) \in A(p)$  and suppose that

$$|f^{(p+1)}(z)| < 2(p!) \quad \text{in } U,$$

then  $f(z)$  is  $p$ -valent in  $U$ .

Applying Schwarz's lemma, it is easily confirmed that Theorem A and the following Theorem A' are equivalent.

Theorem A'. If  $f(z) \in A(1)$  and  $|zf''(z)| < 1$  in  $U$ , then  $f(z)$  is univalent in  $U$ .

We here obtain the following result concerned with Theorem A and Theorem A'.

Theorem 1. Let  $f(z) \in A(1)$  and suppose that

$$(1) \quad \left| \arg \left( zf''(z) + \frac{1}{2} \right) \right| < \pi \quad \text{in } U.$$

Then  $f(z)$  is univalent in  $U$ .

Proof. Let us put  $p(z) = f'(z)$  and

$$\phi(z) = \frac{1 - p(z)}{1 + p(z)}.$$

Then, if there exists a point  $z_0 \in U$  such that

$$\operatorname{Re} p(z) > 0 \quad \text{for } |z| < |z_0|$$

$$\operatorname{Re} p(z_0) = 0,$$

then we have that  $\phi(0) = 0$ ,  $|\phi(z)| < 1$  for  $|z| < |z_0|$  and  $|\phi(z_0)| = 1$ .

Applying the same method as the proof of [3, Theorem 1], we have that  $z_0 p'(z_0)$  is a real number and

$$z_0 p'(z_0) \leq -\frac{1}{2} (1 + |p(z_0)|^2) \leq -\frac{1}{2}.$$

This contradicts (1). Therefore we have

$$\operatorname{Re} f'(z) > 0 \quad \text{in } U.$$

From Noshiro's theorem [1],  $f(z)$  is univalent in  $U$ . This completes our proof.

Concerning a sufficient conditions for the univalence of  $f(z)$ , it is trivial that Theorem 1 is better than Theorem A and Theorem A'.

Theorem 2. Let  $f(z) \in A(p)$  and suppose that

$$\left| \arg \left( z f^{(p+1)}(z) + \frac{p!}{2} \right) \right| < \pi \quad \text{in } U.$$

Then  $f(z)$  is  $p$ -valent in  $U$ .

Proof. Applying the same method as the proof of Theorem 1, we have

$$\operatorname{Re} f^{(p)}(z) > 0 \quad \text{in } U.$$

Then, from Ozaki's theorem [4, Theorem 2],  $f(z)$  is  $p$ -valent in  $U$ .

This completes our proof.

Theorem 2 is also better than Theorem B.

## References

- [1] K. Noshiro: On the theory of schlicht functions. J. Fac. Sci. Hokkaido Univ., (1)2, 129-155 (1934-1935).
- [2] M. Nunokaw, O. Kwon and N. K. Cho: On the multivalent functions, Tsukuba J. Math., (1)15, 141-143 (1991).
- [3] M. Nunokawa: On properties of non-Carathéodory functions, Proc. Japan Acad., 68A(6), 152-153 (1992).
- [4] S. Ozaki: On the theory of multivalent functions, Sci. Rep. Tokyo Bunrika Daigaku A, 2, 167-188 (1935).
- [5] S. Ozaki, I. Ono and T. Umezawa: On a general second order derivative, Sci. Rep. Tokyo Bunrika Daigaku, A, 5, 111-114 (1956).